



DIMENSION 270-G[®]

A GRANULE PREEMERGENCE TURF HERBICIDE

Dimension[®] 270-G is a selective herbicide that provides preemergence control of listed annual grasses, and broadleaf weeds in ornamental landscape areas, established lawns and ornamental turf, including golf course fairways, roughs and tee boxes. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

ACTIVE INGREDIENT:

*Dithiopyr, 3,5-pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S, S-dimethylester

0.27%

INERT INGREDIENTS:

99.73%

TOTAL 100.00%

*This product contains 0.135 pounds of the active ingredient, dithiopyr, per 50 pound bag. Product protected by U.S. Patent No. 4,692,184. Other patents pending.

EPA REG. No. 7001-375

EPA EST. No. 66196-CA-1

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES—Immediately flush with plenty of water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff may adversely affect aquatic organisms. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

GENERAL INFORMATION

This product is a selective herbicide that provides preemergence control of crabgrass and other susceptible annual grasses and broadleaf weeds in established lawns and ornamental turf. This product will not harm nearby established ornamentals when used according to label directions. In New York State this product may only be used by professional applicators at no more than 185 pounds (0.5 pounds active ingredient) per year and is prohibited from use in Nassau and Suffolk Counties.

PRECAUTIONS

Use of this product on golf greens is not recommended. DO NOT apply this product to Colonial bentgrass. For best results, cultural practices that disturb the soil, such as core, spike, hydro-aerification, or verticutting, and top dressing should be done before application of this product. This product should not be applied until turfgrasses have recovered from stresses related to these or other mechanical operations. Irrigation or rainfall within 7 days after application will improve the performance of this product.

Reseeding or Sprigging: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrass. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (slit seeders) designed to give good seed-to-soil contact. This product may be used on seeded, sodded, or sprigged turfgrass that are well-established. Allow the turf to develop a good root system and uniform stand before application.

Use of this product on turfgrass not well-established, weakened by lack of moisture, chemical injury, insects, disease, or temperature extremes that cause stress will increase the potential for injury. Allow turfgrass weakened by these stresses to recover before application of this product. DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. DO NOT allow domestic animals to feed or forage on the treated turf.

This product may be applied as a single application or as a split application in the spring, summer or fall. DO NOT apply more than 185 pounds of this product per acre per application and no more than 555 pounds (185 pounds in New York) of this product per acre per year. Do not apply in Nassau and Suffolk Counties of New York.

TURFGRASS

When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

Cool-Season Grasses: Bentgrass, creeping* (*Agrostis stolonifera*), Bluegrass, Kentucky (*Poa pratensis*), Fescue, fine** (*Festuca rubra*), Fescue, tall (*Festuca arundinacea*), Rye grass, perennial (*Lolium perenne*).

Warm-Season Grasses: Bahiagrass (*Paspalum notatum*), Bermudagrass (*Cynodon dactylon*), Buffalograss (*Buchloe dactyloides*), Carpetgrass (*Axonopus affinis*), Centipedegrass (*Eremochloa ophiuroides*), Kiku-yugrass (*Pennisetum clandestinum*), St. Augustinegrass (*Stenotaphrum secundatum*), Zoysiagrass (*Zoysia japonica*).

Products For: LANDSCAPE • GOLF COURSE • HORTICULTURE • LAWN CARE • SOD PRODUCTION

J.R. Simplot Company • P.O. Box 198, Lathrop, CA 95330 • Toll-Free (800) 992-6066 or (209) 858-2511 • Fax (209) 858-2519 • E-mail: best@simplot.com

Use of this product on certain varieties of creeping bentgrass, such as "Carmen", "Cohansey", or "Seaside" will result in undesirable turf injury. Not all varieties of creeping bentgrass have been tested.

DO NOT apply this product to Colonial bentgrass.

**The following fine fescue varieties have been found to be sensitive to this product: "Atlanta", "Banner", "Beauty", "Bligart", "CF-2", "Enjoy", "HF-93", "Highlight", "Ivalo", "Jamestown", "Koket", "Majenta", "Mary", "Pennlawn", "Tamara", "Tatjana", "Waldorf", and "Waldina"

APPLICATION EQUIPMENT AND INSTRUCTIONS

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Calibrate application equipment prior to usage, and apply this product uniformly over the turfgrass area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous, (see *spreader settings for half rate*). Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

APPLICATION DIRECTIONS

This product provides preemergence control of listed annual grasses and broadleaf weeds when applied prior to their germination in established turfgrass.

CONTROL OF ANNUAL GRASSES AND/OR ANNUAL BROADLEAF WEEDS

This product should be applied prior to the germination of annual grasses and/or broadleaf weeds. Use the higher recommended rates where weed infestations are dense and/or applications are made early and longer residual control is desirable. For preemergence control and/or suppression of annual bluegrass and listed winter annual broadleaf weeds, apply this product prior to their germination and emergence. Use the higher recommended use rates where weed infestations are dense.

Refer to the following table for recommended use rates of this product for the preemergence control of annual grasses and/or broadleaf weeds.

WEEDS CONTROLLED AND USE RATE RECOMMENDATIONS			
*USE 100 TO 150 POUNDS/ACRE = 2.3 TO 3.4 POUNDS/1000 SQ.FT.			
FOR: Crabgrass; (Large, Smooth, Southern), Foxtail; (Green, Yellow)			
*USE 150 TO 185 POUNDS/ACRE = 3.4 TO 4.2 POUNDS/1000 SQ.FT.			
FOR:	Barnyardgrass	Lespedeza, common	Ryegrass, Annual
	Chickweed	Kikuyugrass	Perennial
	Clover, hop	Oxalis,	Smutgrass
	Crabgrass, tropical	Buttercup	Speedwell, corn
	Crowfootgrass	Creeping Woodsorrel	Spurge, Prostrate
	Dichondra, Carolina	Yellow Woodsorrel	Spotted
		Pineappleweed	
**USE 185 POUNDS/ACRE = 4.2 POUNDS/1000 SQ.FT.			
FOR:	Bluegrass, annual	Henbit	
	Bittercress	Parsley-piert	
	Geranium, Carolina	Pineappleweed	
	Chickweed	Purslane, common	
	Goosegrass	Shepardspurse	
*The higher use rates are recommended where weed infestations are heavy and/or dense.			
**Use of this product at this rate will provide only partial control and/or suppression of listed weeds.			

SPREADER SETTINGS

Pounds / 1000 sq.ft.	Half Rate Application			Full Rate Application		
	1.15 lbs.	1.7 lbs.	2.1 lbs.	2.3 lbs.	3.4 lbs.	4.2 lbs.
BEST® PRO	3.5	3.75	3-7/8 A	4A	4-3/8A	4-1/2A
BEST® 34, 44, 64	2.8	3	3.1	3.2	3.7	3.9
SCOTTS R8A	G Cone 8	G-1/2 Cone 8	H Cone 8	H-1/2 Cone 8	I-1/2 Cone 8	J Cone 8
LELY TYPE W	3	3.5	4	4.5	5	5.5
EARTHWAY	9-1/2	10	10-1/2	11	12	12-1/2

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: Store this product in its original container only. In case of spill, sweep up material and dispose of material according to label directions.

DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, State or Local procedures. Do not reuse emptied

GENERAL INFORMATION - LANDSCAPE ORNAMENTALS

DIMENSION® 270-G provides preemergence control of listed annual grass and broadleaf weeds in landscape ornamentals. It may be applied to soil surfaces where ornamental plants are grown for aesthetic purposes in ornamental gardens, parks, golf courses and residential properties. Ornamental plants on this label have shown tolerance to DIMENSION® 270-G when applied as directed. It is impossible, however, to evaluate tolerance for all ornamental plant species or varieties or under all possible growing conditions. Under your growing conditions, a limited area involving only a few plants should be treated for tolerance evaluation prior to large-scale applications.

USE RATES: Apply DIMENSION® 270-G with a properly calibrated spreader which will assure accurate, even particle distribution. This product may be applied as a single application or as a split application. Refer to the following table for product recommendations when treating a small area.

POUNDS PRODUCT/ACRE	OUNCES PRODUCT/1000 SQ.FT.	OUNCES PRODUCT/100 SQ.FT.
185	68	6.8 (193 grams)

APPLICATION TIMING AND RECOMMENDATIONS: DIMENSION® 270-G is a preemergence herbicide that will not control established weeds. Make applications prior to weed seed germination or to bare ground. The best weed control is obtained when applied to soil that is free of clods, weeds and debris such as leaves. Prior to making an application, control existing vegetation by hand weeding, cultivation or with a postemergence herbicide. Once an application of this product has been made, do not disturb the soil surface as the herbicide barrier will be broken.

PRECAUTIONS:

- Only apply this product to established ornamentals.
- Do not apply this product directly to bare root of ornamental plants as injury may result.
- Do not incorporate this product into soil. Dilution of active ingredient and possible injury to plant roots may occur.
- Do not apply to soil around ornamental plants that are under stress due to drought, flooding, excessive fertilizer or soil salts, wind injury, hail, frost damage, winter injury, injury from previously applied pesticides or injury due to insects, nematodes or disease.
- To avoid foliar damage to ornamentals, irrigate immediately after application to remove herbicide particles from the foliage.
- Do not apply under conditions that would affect product distribution. Wind could affect distribution and move product onto sensitive species. Uneven product distribution will result in uneven weed control.
- Do not apply this product on grasses grown for seed
- Do not graze livestock or feed forage cut from areas treated with this product.

TOLERANT ORNAMENTALS: When applied as directed under the conditions described on this label, ornamentals listed below have shown tolerance in field trials. However, this product has not been tested on all cultivars of each species or under all possible growing conditions. Follow directions given above to determine plant tolerance under your growing conditions prior to large scale use.

Abelia Dwarf (Abelia X grandiflora) Nana; **Ajuga** (Ajuga reptans, Ajunga genevensis) Bronze, Bronze Beauty; **Almond, Flowering** (Prunus glandulosa); **Apple*** (Malus pumila); **Arborvitae** (Thuja occidentalis) Nigra, Pyramidalis, Smaragd, Techny, Woodwardii; **Arborvitae, Golden** (Thuja orientalis); **Aster, Chinese** (Callistephus chinensis) Dwarf Queen; **Ash, Green** (Fraxinus pennsylvanica); **Ash, Mountain** (Sorbus aucuparia); **Ash, Purple** (Fraxinus americana); **Azalea** (Rhododendron spp.) Brilliant, Buccaneer, Carror, Chimes (Belgian), Exbury, Fashion, Hardijzer Beauty, Hershey Red, Higasa, Hinocrimson, Holland (Hybrid) Marion Lee, Northern Lights, Orange Cup, Orchid Lights, Snow, Southern Charm; **Azalea, Flame** (Rhododendron calendulaceum); **Azalea, Kirishima; Bamboo, Heavenly; Barberry** (Berberis thunbergii) Aurea, Dwarf Pigmy, Green, Kobold, Pygmy Red, Rose Glow; **Barberry, Purple**, Atropurpurea; **Bearberry (common)** (Arctostaphylos uva-ursi) Massachusetts; **Birch, River** (Betula nigra); **Blueberry*** (Vaccinium spp.) Bluecrop, Blue Jay, Jersey, North Blue, Northland; **Bottlebrush** (Callistemon citrinus); **Boxwood, Japanese**, Japonica; **Boxwood, Weller** (Buxus sempervirens); **Broom** (Cytisus spp.) Moonlight; **Broom** (Genista pilosa) Vancouver Gold; **Bugle Carpet; Carex, Variegated** (Carex); **Camellia** (Camellia japonica, Camellia sasangua) Debutante, Mathotiana Supreme, Chansonette; **Cedar, Red** (Juniperus virginiana); **Cockscomb, Plumosa** (Celosia cristata) Red Kewpie; **Coleus** (Coleus blumei); **Copper leaf** (Acalypha wilkesiana); **Cotoneaster** (Cotoneaster apiculatus); **Cycads** (Cycads revoluta); **Coyotebrush** (Baccharis pelularis); **Cypress, Bald** (Taxodium distichum); **Cypress, Italian** (Cupressus sempervirens) Glauca; **Cypress, Leyland** (Cupressocyparis leylandii); **Cypress, Japanese False** (Chamaecyparis obtusa) Gracilis; **Daffodil** (Narcissus spp.) King Alfred; **Daylily** (Hemerocallis spp.) Aztec Gold, Bright Yellow (Hybrid), Single Gold (Evergreen), Wilson's Yellow; **Dogwood** (Cornus florida); **Dogwood, American** (Cornus sericea) Flava; **Douglas Fir** (Pseudotsuga menziesii); **Dusty Miller** (Senecio cineraria); **Elm, Drake** (Ulmus parvifolia); **Euonymus** (Euonymus fortunei) Argenteo-variegata, Auereu-marginata, Colorata, Emerald Gaiety, Emerald 'N Gold, Gold Edge, Gold Princess, Silver King, Tricolor, Vegetus; **Fan Palm, European** (Chamaerops humilis); **Fan Palm, Mexican** (Washingtonia robusta); **Fescue** (Festuca glauca); **Fetterbush** (Leucothoe fontiana) Rainbow; **Ficus** (Ficus retusa) Nitidia; **Forsythia** (Forsythia X intermedia) Arnold Dwarf, Bronxensis Dwarf, Lynwood Gold, Meadowlark, (Spring Ligo) Weeping; **Fir Fraser** (Abies fraseri); **Gardenia** (Gardenia jasminoides) Mystery, Radicans; **Geranium** (Pelargonium X hortorum); **Gum** (Eucalyptus citriodora); **Hawthorn** (Crataegus spp.) Cockspur White, Crimson Cloud, Enchantress, Jack Evans, Washington White; **Heather, Twisted** (Erica cinerea) Mediterranean Pink; **Hibiscus** (Hibiscus spp.) Brilliant, Hula Girl, Blue Bird; **Hemlock, Canada** (Tsuga canadensis); **Holly** (Ilex spp.) Blue Boy, Blue Girl, (Ilex X meserveae) Burfordii, China Girl, Compacta, Forsteri, Hellerie, Japanese Northern Beauty, Needlepoint, Nellie R. Stevens, Savanah, (Ilex X attenuata); **Holly, Chinese** (Ilex cornuta); **Holly, Japanese** (Ilex crenata); **Holly, Yaupon** (Ilex vomitoria); **Honeysuckle** (Lonicera japonica) Clavey's Dwarf, Halliana, Tatarian Canadian White, Zebelli Red Hosta; **Hosta** (Hosta spp.) Albo Marginata (Hosta lancifolia); Ice Plant (Carpobrotus edulis); Impatiens (Impatiens wallerana); Iris (Iris spp.) Wedgewood, Dwarf Blue; **Ivy, English** (Hedera helix) Bulgaria, Thordale; **Jasmine, Asian** (Trachelopermum asiaticum); **Juniper** (Juniperus spp.) Arcadia, Armstrong, Bar Harbor--(Juniperus horizontalis) Blue Chip, Blue Pacific, Blue Rug--(Juniperus horizontalis) Blue Star, Broadmoor, Buffalo, Calgary Carpet, Emerald Sea, Emerald Spreader, Endora Compacta, Fruitlandi, Green, Gold Tip, Hetzi, Hughes, Manhattan Blue, Parsoni, Pfitzeriana, Plumosa, Prince of Wales, Procmbers Dwarf, San Jose, Sargent Blue, Sargent Sea, Scandia, Scopolorum Moonglow, Scopolorum Skyrocket, Spartan--(Juniperus chinensis) Tamariscifolia--(Juniperus sabina) Weberi, Youngstown, Yukon Belle; **King Palm** (Archontophoenix cunninghamiana); **Laurel, Australian** (Pittosporum tobira); **Laurel, Mountain** (Kalmia latifolia); **Leucothoe** (Leucothoe fontanesiana); **Lily, African** (Agapanthus africanus) Albus, Peter Pan; **Lily, African Blue; Lily of the Valley** (Pieris japonica) Mt. Fire; **Lilyturf** (Liriope muscar) Evergreen Giant, Lilac Beauty, Majestic, Monroe White, Variegata; **Ligustrum, Japanese** (Ligustrum japonicum); **Liriope, Green** (Liriope spicata); **Maple, Japanese** (Acer japonicum); **Maple, Norway** (Acer platanoides); **Maple, Red** (Acer rubrum); **Maple, Silver** (Acer saccharinum); **Maple Sugar** (Acer saccharum); **Marigold** (Tagetes patula) Honeycomb, Wheeler's Dwarf, Variegata; **Mock Orange*** (Philadelphus spp.) Golden, Snowflake Double White; **Monpdo Grass** (Ophiopogon japonicus); **Miss Rose** (Portulaca grandiflora) Sunnyside; **Myrtle, Crane** (Lagerstemia indica) Faurei, Langer, Muskogee, Standard Pink; **Myrtle Wax** (Myrica cerifer); **Nadina** (Nadina domestica) Compacta, Nana; **Oleander** (Nerium oleander) Hardy Red, Petite Pink, Sister Agnes; **Oak, Laurel** (Quercus laurifolia); **Oak, Pin** (Quercus palustris); **Oak, Red** (Quercus rubra); **Oak, Southern L** (Quercus virginiana); **Oak, Willow** (Quercus phellos); **Osteospermum** (Osteospermum fruticosum) Wirilrig; **Palm, Bangalow; Pampas Grass** (Cortadaria setoana); **Paper Flower** (Bougainvillea glabra) Barbara Karst; **Pachysandra** (Pachysandra terminalis); **Peach*** (Prunus persica); **Periwinkle, Dwarf** (Vinca minor); **Petunia** (Petunia X hybrida) Picotti; **Photinia, Red Tip** (Photinia X fraseri); **Pieris** (Pieris japonica); **Pine, Australian** (Pinus nigra); **Pine, Japanese Black** (Pinus thumbergia); **Pine, Loblolly** (Pinus taeda); **Pine, Longleaf** (Pinus palustris); **Pine, Mugho** (Pinus mugho); **Pine, Scotch** (Pinus sylvestris); **Pine, Slash** (Pinus elliotii); **Pine, Swiss Mt.** (Pinus mugho); **Pine, Virginia** (Pinus virginiana); **Pine, White** (Pinus strobus); **Pineapple, Guava** (Feijoaellowaiiana); **Pittosporum, Japan; Potentilla** (Potentilla fruticosa) Abbotswood; **Privet** (Ligustrum japonicum) Golden Vicary, Regal, Texanum, Yellow Tipped, Wax; **Privet, Glossy** (Ligustrum lucidum); **Pyracantha** (Pyracantha koidzumii) Gnome, Lalanderi, Victory; **Queen Palm** (Arecastrum rammanzoffianum); **Quince, Japanese***; **Rhododendron** (Rhododendron spp.) Album, Cunningham White, Fashion, Hardy, PJM, Purple Gem, Silvery Pink; **Rhododendron, Carolina** (Rhododendron carolinianum); **Rhododendron, Catawba** (Rhododendron catawbiense); **Rodie Max (Rosebay)** (Rhododendron maximum); **Rose** (Sosa banksiaae) Luta; **Rosemary*** (Rosmarinus officinalis); **Rosemary, Bog** (Andromeda polifolia) Nana; **Salvia** (Salvia farinacea) Rhea; **Sedum** (Sedum spurius) Dragon Blood Red, Red Carpet, Yellow; **Spiraea** (Astilbe X arendsi) Fanall; **Spiraea** (Spiraea spp.) Anthony Waterer Red, Dolchica, Froebel Pink, Goldenflame Red, Snowmound White, Van Houtte White; **Spiraea, Garland** (Spiraea spp.); **Spruce, Black Hills** (Picea glauca); **Spruce, Colorado Blue** (Picea pungens); **Spruce, Norway** (Picea abies); **Spruce, White** (Picea glauca) Conica; **Sourwood** (Oxydendrum arboreum); **Sweetflag, Grassyy-Lc; Sweetgum** (Liquidambar styraciflua); **Sycamore** (Platanus occidentalis); **Tree Fern (Tiki Fern)** (Asparagus virgatus); **Trumpet Flower, Evening** (Gelsemium sempervirens); **Tulip** (Tulip, spp.) Appleddown; **Verbana, Shrub** (Lantana sellowiana); **Vervain** (Verbena spp.) St. Paul; **Viburnum** (Viburnum spp.) American Cranberry Bush, Arrowwood, Common Snowball, European Cranberry Bush, Linden, Mohican, Wright; **Windmill Palm** (Trachycarpus fortunei); **Xylosma** (Xylosma congestum); **Yaupon** (Ilex vomitoria) Dwarf; **Yew** (Taxus cuspidata, Taxus X media) Denisiformis.

*Ornamental species only: DO NOT USE ON FOOD PRODUCING TREES AND ORNAMENTALS.

WEEDS CONTROLLED: Used as directed, DIMENSION® 270-G will control or suppress the following annual grass and broadleaf weeds when applied prior to their emergence. This product will not control established grasses, except for crabgrass in early stages of development, and broadleaf weeds, therefore, the area to be treated should be free of weeds prior to application.

Grasses: barley (Hordeum spp.); barnyardgrass (Echinochloa crus-galli); bluegrass, annual (Poa annua); brome (Bromus spp.); crabgrass, large (Digitaria sanguinalis); crabgrass, smooth (Digitaria ischaemum); crabgrass, Southern (Digitaria ciliaris) crowfootgrass (Dactyloctenium aegyptium); dallisgrass (seedling) (Paspalum dilatatum); goosegrass (Eleusine indica); foxtail, green (Setaria verticillata); foxtail, yellow (Setaria faberi); kikuyugrass (Pennisetum clandestinum) oats, wild (Avena fatua); ryegrass (annual & perennial) (Lolium spp.); smutgrass (Sporobolus poireti).

Broadleaf Weeds: bittercress (Cardamine spp.); carpetweed (Mollugo verticillata); chickweed (Stellaria spp.); geranium, Carolina (Geranium carolinianum); henbit (Lamium spp.); knotweed, prostrate (Polygonum saviculare); lespedeza, common (Lespedeza striata); marestail (Hippuris vulgaris); medic, black (Medicago lupulina); mustard (Brassica spp.); oxalis, buttercup (Oxalis pes-caprae); pineappleweed (Matricaria matricarioides); pigweed, redroot (Amaranthus retroflexus); parsley-piert (Alchemilla arvensis); purslane, common (Portulaca oleracea); rocket, London (Sisymbrium irio); shepherdspurse (Capsella bursa-pastoris); speedwell, corn (Veronica arvensis); spurge, garden (Euphorbia hirta); spurge, prostrate (Euphorbia humistrata); spurge, spotted (Euphorbia maculata); woodsorrel, creeping (Oxalis corniculata); woodsorrel, yellow (Oxalis stricta).

APPLICATION EQUIPMENT AND INSTRUCTIONS: Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the area to be treated. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

50 LB. BAG (22.68 Kg.)

Treats up to 21,780 Sq.Ft. for Crabgrass Control

DISCLAIMER OF WARRANTIES: Seller warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation are beyond the control of seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of seller's liability. No one is authorized to make any other warranty, guarantee or directions concerning this product, and not such warranties, guarantees or directions shall be valid or binding upon seller.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-jr.com>